



A FLY AND LICE SPRAY-ON FOR SHEEP

ZENITH Spray-on is an IGR spray-on for rapid maggot kill, the prevention of flystrike and control of lice



- Diflubenzuron (IGR) pour-on for sheep
- Treatment of active blowfly strike
- Prevention of blowfly strike and control of lice
- 7 day meat withholding period
- Provides up to 12 weeks fly control
- Provides 20 weeks lice control

Registered for:

The treatment of active blowfly strike, and prevention of blowfly strike with up to 12 weeks control. Also for the control of lice with 20 weeks control.

ZENITH Spray-on is an insect growth regulator (IGR) fly and lice pour-on for sheep, containing the active diflubenzuron. Diflubenzuron acts by interfering with the moulting and growth process causing death of blowfly larvae and lice nymphs. ZENITH Spray-on also contains hydrocarbon liquid, which kills maggots on contact to treat active blow-fly strike.

ZENITH Spray-on treats both fly and lice, so is ideal throughout the fly season for the prevention of flystrike, and to ensure the control of lice before going into the colder months, when lice can become a problem.

The added convenience of treating active flystrike means ZENITH Spray-on is especially useful to have on-hand over the fly season. And our 900mL spray bottle fits nicely in the farmbike tool box.

More great reasons to buy ZENITH Spray-on:

- **FREE GUN** Receive a free applicator gun when you buy 2 x 20L drums of ZENITH Spray-on
- **PRIORITY PARTNERSHIP POINTS** Earn 250 Priority Partnership Points for every 20L and 60 Priority Partnership Points for every 5L ZENITH Spray-on drum that you buy.

Withholding times:

Meat: 7 days

Milk: 35 days

Wool Harvest Interval: 60 days

Application:

Apply ZENITH Spray-on using a COOPERS T-bar gun. Apply in a continuous band down the midline from between the ears (lice) or neck to the tail. For fly, also apply to the crutch area.

Dose rate:


For treatment of active strike, apply directly to maggots in fly-struck area. Clip wool from the affected area (do not shear) to prevent discharge build-up in the fleece.

Fly and lice dose rates overleaf.




Dose rate: Fly


Class/Liveweight	Dose	Apply To	Doses Per Pack		
			900mL	5L	20L
Docking/Tailing	10-15mL (1x10-15mL)	Crutch	60	333	1,333
Lambs <20kg	20mL (2x10mL)	Crutch/Back	45	250	1,000
Lambs >20kg Hoggets & Ewes	30mL (2x15mL)	Crutch/Back	30	166	666
Heavy Ewes & Rams	45mL (3x15mL)	Crutch/Back	20	111	444




Docking



0-20kg



20kg+



Heavy ewes
and rams

Dose rate: Lice

Liveweight	Dose	Apply To	Doses Per Pack		
			900mL	5L	20L
<30kg	15mL	Neck & Backline	60	333	1,333
30-55kg	20mL	Neck & Backline	45	250	1,000
56-75kg	25mL	Neck & Backline	36	200	800

